



SAVE THE DATE

3RD PREMICOSE SYMPOSIUM

PDA Closure in Pre-term Infants

DECEMBER 2ND 2022, SPAIN

PANEL 1 – After 10 years where are we now? Technical variations. Different ways to do it

PANEL 2 – What to do to limit complications? How to deal with complications?
Establishing a Decision-Making Protocol

SHORT TOPICS AND CASES PRESENTATIONS – It's still not clear to me to close the PDA

PANEL 3 – What evidence do we have on early effect of transcatheter PDA closure?
What do we know when comparing with surgery?

PANEL 4 – What do we know on midterm evolution? What can we improve?

PANEL 5 – Some questions about the percutaneous closure of the preterm PDA
“not everything is clear”

CAUTION: This product is intended for use by or under the direction of a physician. Prior to use, reference the Instructions for Use, inside the product carton (when available) or at eifu.abbottvascular.com or at medical.abbott/manuals for more detailed information on Indications, Contraindications, Warnings, Precautions and Adverse Events. Device contraindicated for patients with a weight < 700 grams at time of the procedure. Illustrations are artist's representations only and should not be considered as engineering drawings or photographs. Photo(s) on file at Abbott. Information contained herein for DISTRIBUTION in Europe, Middle East and Africa ONLY. Please check the regulatory status of the device before distribution in areas where CE marking is not the regulation in force.

For more information, visit our web site at www.abbott.com

Abbott Vascular International BVBA

Park Lane, Culliganlaan 2B, B-1831 Diegem, Belgium, Tel: +32 2 714 14 11

© 2022 Abbott. All rights reserved. 9-EH-8-13503-01 07-2022

